

Meson decay in the Fock-Tani Formalism

D.T. da Silva*, M. L. L. da Silva*, D. Hadjimichef^{†,*} and C. A. Z. Vasconcellos*

**Instituto de Física, Universidade Federal do Rio Grande do Sul, Caixa Postal 15051, CEP 91501-970, Porto Alegre, RS, Brazil.*

[†]Departamento de Física, Instituto de Física e Matemática, Universidade Federal de Pelotas, Campus Universitário, CEP 96010-900, Pelotas, RS, Brazil.

Abstract.

The Fock-Tani formalism is a first principle method to obtain effective interactions from microscopic Hamiltonians. Usually this formalism was applied to scattering, here we introduced it to calculate partial decay widths for mesons.

Keywords: meson decay; Fock-Tani formalism

PACS: 13.25.Jx ; 12.39.Pn

INTRODUCTION

In the Fock-Tani representation [1] one starts with the Fock representation of the system using field operators of elementary constituents which satisfy canonical (anti) commutation relations. Composite-particle field operators are linear combinations of the elementary-particle operators and do not generally satisfy canonical (anti) commutation relations. "Ideal" field operators acting on an enlarged Fock space are then introduced in close correspondence with the composite ones. The enlarged Fock space is a graded direct product of the original Fock space and an "ideal state space". The ideal operators correspond to particles with the same quantum numbers of the composites; however, they satisfy by definition canonical (anti)commutation relations. Next, a given unitary transformation, which transforms the single composite states into single ideal states, is introduced. When the transformation acts on operators in the subspace of the enlarged Fock space which contains no ideal particles, the transformed operators explicitly express the interactions of composites and constituents. Application of the unitary operator on the microscopic Hamiltonian, or on other hermitian operators expressed in terms of the elementary constituent field operators, gives equivalent operators which contain the ideal field operators. In the present we apply the formalism to meson decay into two-mesons. In particular the example system is $\rho^+ \rightarrow \pi^+ \pi^0$.

THE FOCK-TANI FORMALISM

Now let us to apply the Fock-Tani formalism in the microscopic Hamiltonian to obtain an effective Hamiltonian. In the Fock-Tani formalism we can write the meson creation

operators in the following form

$$M_{\alpha}^{\dagger} = \Phi_{\alpha}^{\mu\nu} q_{\mu}^{\dagger} \bar{q}_{\nu}^{\dagger}.$$

where $\Phi_{\alpha}^{\mu\nu}$ is the bound-state wave-functions for two-quarks respectively. The operators of the quark and antiquark obey the following anticommutation relations

$$\begin{aligned} \{q_{\mu}, q_{\nu}\} &= \{\bar{q}_{\mu}, \bar{q}_{\nu}\} = \{q_{\mu}, \bar{q}_{\nu}\} = \{q_{\mu}, \bar{q}_{\nu}^{\dagger}\} = 0 \\ \{q_{\mu}, q_{\nu}^{\dagger}\} &= \{\bar{q}_{\mu}, \bar{q}_{\nu}^{\dagger}\} = \delta_{\mu\nu}. \end{aligned}$$

The composite meson operators satisfy non-canonical commutation relations

$$[M_{\alpha}, M_{\beta}] = 0; [M_{\alpha}, M_{\beta}^{\dagger}] = \delta_{\alpha\beta} - \Delta_{\alpha\beta}$$

where

$$\Delta_{\alpha\beta} = \Phi_{\alpha}^{*\mu\gamma} \Phi_{\beta}^{\gamma\rho} q_{\rho}^{\dagger} q_{\mu} + \Phi_{\alpha}^{*\mu\gamma} \Phi_{\beta}^{\gamma\rho} \bar{q}_{\rho}^{\dagger} \bar{q}_{\mu}.$$

The idea of the Fock-Tani formalism is to make a representation change, of form that the composite particles operators are described by operators that satisfy canonical commutation relations. i.e., which obey canonical relations

$$[m_{\alpha}, m_{\beta}] = 0; [m_{\alpha}, m_{\beta}^{\dagger}] = \delta_{\alpha\beta}.$$

where $m_{\alpha}^{\dagger}$ is the operator of the “ideal particle” creation. This way one can transform the composite state $|\alpha\rangle$ into an ideal state $|\alpha\rangle$, in the meson case for example we have

$$U^{-1} M_{\alpha}^{\dagger} |0\rangle = m_{\alpha}^{\dagger} |0\rangle = |\alpha\rangle$$

where $U = \exp(tF)$ and F is the generator of the meson transformation given by

$$F = \sum_{\alpha} m_{\alpha}^{\dagger} \tilde{M}_{\alpha} - \tilde{M}_{\alpha}^{\dagger} m_{\alpha}$$

with

$$\tilde{M}_{\alpha} = M_{\alpha} + \frac{1}{2} \Delta_{\alpha\beta} M_{\beta} + \frac{1}{2} M_{\beta}^{\dagger} [\Delta_{\beta\gamma}, M_{\alpha}] M_{\gamma}.$$

A set of Heisenberg-like equations for the basic operators m, M, q

$$\frac{dm_{\alpha}(t)}{dt} = [m_{\alpha}(t), F] = \tilde{M}_{\alpha}(t); \quad \frac{d\tilde{M}_{\alpha}(t)}{dt} = [\tilde{M}_{\alpha}(t), F] = -m_{\alpha}(t).$$

The simplicity of these equations are not present in the equations for q

$$\frac{dq_{\mu}(t)}{dt} = [q_{\mu}(t), F] = -\Phi_{\alpha}^{\mu\nu} \bar{q}_{\nu}^{\dagger}(t) m_{\alpha}(t)$$

and

$$\frac{d\bar{q}_v(t)}{dt} = [\bar{q}_v(t), F] = \Phi_\alpha^{\mu\nu} q_\mu^\dagger(t) m_\alpha(t).$$

The solution for these equations can be found order by order in the wavefunctions. For zero order one has $q_\mu^{(0)}(t) = q_\mu$, $\bar{q}_v^{(0)}(t) = \bar{q}_v$, $m_\alpha^{(0)}(t) = M_\alpha \sin t + m_\alpha \cos t$, $M_\beta^{(0)}(t) = M_\beta \cos t - m_\beta \sin t$. In the first order $m_\alpha^{(1)} = 0$, $M_\beta^{(1)} = 0$ and $q_\mu^{(1)}(t) = -\Phi_\alpha^{\mu\nu_1} \bar{q}_{\nu_1}^\dagger (M_\alpha - m_\alpha)$; $\bar{q}_v^{(1)}(t) = \Phi_\alpha^{\mu_1\nu} q_{\mu_1}^\dagger (M_\alpha - m_\alpha)$. for these calculations, we are using $t = -\frac{\pi}{2}$ Consequently, for second and third order, one obtains

$$\begin{aligned} q_\mu^{(2)}(t) &= \frac{1}{2} \Phi_\alpha^{*\mu_2\nu_1} \Phi_\beta^{\mu\nu_1} [-m_\alpha^\dagger m_\beta - M_\alpha^\dagger M_\beta + M_\alpha^\dagger m_\beta] q_{\mu_2} \\ \bar{q}_v^{(2)}(t) &= \frac{1}{2} \Phi_\alpha^{*\mu_1\nu_2} \Phi_\beta^{\mu_1\nu} [-m_\alpha^\dagger m_\beta - M_\alpha^\dagger M_\beta + M_\alpha^\dagger m_\beta] \bar{q}_{\nu_2} \end{aligned}$$

and

$$\begin{aligned} q_\mu^{(3)}(t) &= \frac{1}{2} \Phi_\alpha^{*\rho\sigma} \Phi_\beta^{\mu\sigma} \Phi_\gamma^{\rho\sigma_1} \bar{q}_{\sigma_1}^\dagger [-m_\alpha^\dagger m_\beta m_\gamma - M_\alpha^\dagger m_\beta M_\gamma + m_\alpha^\dagger m_\beta M_\gamma \\ &\quad + M_\alpha^\dagger M_\beta M_\gamma] + \frac{1}{2} \Phi_\alpha^{*\rho\sigma} \Phi_\alpha^{\mu\nu_1} \Phi_\beta^{\rho\sigma_1} \bar{q}_{\nu_1}^\dagger \bar{q}_{\sigma_1}^\dagger \bar{q}_\sigma [M_\beta + m_\beta] \\ &\quad + \frac{1}{2} \Phi_\alpha^{*\rho\sigma} \Phi_\alpha^{\mu\nu_1} \Phi_\beta^{\rho_1\sigma} \bar{q}_{\nu_1}^\dagger q_{\rho_1}^\dagger q_\rho [M_\beta + m_\beta] \\ \bar{q}_v^{(3)}(t) &= \frac{1}{2} \Phi_\alpha^{*\rho\sigma} \Phi_\beta^{\rho\nu} \Phi_\gamma^{\rho_1\sigma} q_{\rho_1}^\dagger [-m_\alpha^\dagger m_\beta m_\gamma - M_\alpha^\dagger m_\beta M_\gamma + m_\alpha^\dagger m_\beta M_\gamma \\ &\quad + M_\alpha^\dagger M_\beta M_\gamma] + \frac{1}{2} \Phi_\alpha^{*\rho\sigma} \Phi_\alpha^{\mu_1\nu} \Phi_\beta^{\rho\sigma_1} q_{\mu_1}^\dagger \bar{q}_{\sigma_1}^\dagger \bar{q}_\sigma [M_\beta + m_\beta] \\ &\quad + \frac{1}{2} \Phi_\alpha^{*\rho\sigma} \Phi_\alpha^{\mu_1\nu} \Phi_\beta^{\rho_1\sigma} q_{\mu_1}^\dagger q_{\rho_1}^\dagger q_\rho [M_\beta + m_\beta] \end{aligned}$$

THE MICROSCOPIC MODEL

In this model we use the following Hamiltonian inspired in the 3P_0 approach

$$H = g \int d^3x \Psi^\dagger(\vec{x}) \gamma^0 \Psi(\vec{x}).$$

Where the quark field is respectively

$$\Psi(\vec{x}) = \sum_s \int \frac{d^3p}{(2\pi)^{3/2}} [u(\vec{p}, s) b(\vec{p}, s) + v(-\vec{p}, s) d^\dagger(-\vec{p}, s)] e^{i\vec{p} \cdot \vec{x}}.$$

and $\gamma = \frac{g}{2m_q}$ where m_q is the mass of both produced quarks. After momentum the expansion of the quark field and retaining only the $b^\dagger d^\dagger$ yields

$$H_I = g \sum_{ss'} \int d^3p d^3p' \delta(\vec{p} + \vec{p}') u_{s'}^\dagger(\vec{p}') \gamma^0 v_s(\vec{p}) b_{s'}^\dagger(\vec{p}') d_s^\dagger(\vec{p}).$$

Introducing the following notation $b \rightarrow q; d \rightarrow \bar{q}; \mu = (\vec{p}', s'); \nu = (\vec{p}, s)$ and

$$V_{\mu\nu} = g \sum_{ss'} \int d^3p d^3p' \delta(\vec{p} + \vec{p}') u_{s'}^\dagger(\vec{p}') \gamma^0 v_s(\vec{p}).$$

The Hamiltonian now is reduced to a compact form, $H_I = V_{\mu\nu} q_\mu^\dagger \bar{q}_\nu^\dagger$ where sum (integration) is implied over repeated indexes. Applying the Fock-Tani transformation to H_I one obtains the effective Hamiltonian $H_{IT} = U^{-1} H_I U$. For a $q\bar{q}$ meson A to decay to mesons $B + C$ we must have $(q\bar{q})_A \rightarrow (q\bar{q})_B + (q\bar{q})_C$. The transformed operators that shall contribute to $U^{-1} H_I U$ are

$$H_m = V_{\mu\nu} q_\mu^{\dagger(3)} q_\nu^{\dagger(0)} + V_{\mu\nu} q_\mu^{\dagger(1)} q_\nu^{\dagger(2)}$$

Which results in the effective meson decay Hamiltonian after some manipulation

$$H_m = -\Phi_\alpha^{*\mu\lambda} \Phi_\gamma^{*\rho\sigma} \Phi_\beta^{\rho\lambda} V_{\mu\nu} m_\alpha^\dagger m_\gamma^\dagger m_\beta.$$

We now consider the transition $m_\beta \rightarrow m_\alpha^\dagger m_\gamma$, where $|i\rangle = m_\beta^\dagger |0\rangle$ and $|f\rangle = m_\alpha^\dagger m_\gamma^\dagger |0\rangle$. Finally we find

$$\langle f | H_m | i \rangle = -\Phi_\alpha^{*\rho\nu} \Phi_\beta^{\rho\lambda} \Phi_\gamma^{*\mu\lambda} V_{\mu\nu} - \Phi_\alpha^{*\mu\lambda} \Phi_\beta^{\rho\lambda} \Phi_\gamma^{*\rho\nu} V_{\mu\nu}$$

The meson wave function is defined as

$$\Phi_\alpha^{\mu\nu} = \chi_{S_\alpha, M_\alpha^S}^{S_1 S_2} f_{f_\alpha, M_\alpha^f}^{f_1 f_2} C_{C_\alpha}^{C_1 C_2} \Phi_{\vec{P}_\alpha}^{\vec{p}_i \vec{p}_j}$$

where χ is spin; f is flavor and C are color coefficients. The spatial parte is given by

$$\Phi_{\vec{P}_\alpha}^{\vec{p}_i \vec{p}_j} = \delta^3(\vec{P}_\alpha - \vec{p}_i - \vec{p}_j) \left(\frac{1}{\pi b^2} \right)^{3/4} e^{-\frac{1}{8b^2}(\vec{p}_i - \vec{p}_j)^2}$$

We analyze the particular case $\rho^+(+\hat{z}) \rightarrow \pi^+ \pi^-$ decay. For this decay have that the flavor part is given by $|\rho^+\rangle = |\pi^+\rangle = -|ud\rangle$; $|\pi^0\rangle = \frac{1}{\sqrt{2}}[|u\bar{u}\rangle - |d\bar{d}\rangle]$ and the spin part is given by $|\rho(+\hat{z})\rangle = |\uparrow\uparrow\rangle$; $|\pi\rangle = \frac{1}{\sqrt{2}}[|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle]$. The transition matrix element is given by

$$\langle f | H_{IT} | i \rangle = \delta^3(\vec{P}_\beta - \vec{P}_\alpha - \vec{P}_\gamma) h_{fi}$$

with

$$h_{fi} = \frac{2^3}{3^3} \gamma \left(\frac{1}{\pi^{3/4} b^{1/2}} \right) \frac{P}{b} e^{-\frac{P^2}{12b^2}}$$

This h_{fi} decay amplitude can be combined with relativistic phase space to give the differential decay rate, which is [2]

$$\frac{d\Gamma_{A \rightarrow BC}}{d\Omega} = 2\pi \frac{PE_B E_C}{M_A} |h_{fi}|^2$$




FIGURE 1. Γ_{theor} for $\rho \rightarrow \pi\pi$; $\Gamma_{\text{exp}} = 151 \text{ MeV}$ is obtained for $\beta = 0.397 \text{ GeV}$

resulting

$$\Gamma_{A \rightarrow BC} = \sqrt{\pi} \frac{2^7}{3^6} \left(\frac{g}{2m_q} \right)^2 M_\rho \left(\frac{P}{b} \right)^3 e^{-\frac{P^2}{6b^2}}.$$

CONCLUSIONS

The Fock-Tani formalism is proven appropriate not only for hadron scattering but for decay. The example decay process $\rho^+ \rightarrow \pi^+ \pi^0$ in the Fock-Tani formalism reproduces the predictions in the 3P_0 model. The same procedure can be used for $f_0(M)$ and for heavier scalar mesons and compared with similar calculations with glue content.

ACKNOWLEDGMENTS

The authors acknowledges support from the Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul - FAPERGS. D.T.S. acknowledges support from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - CAPES. M.L.L.S. acknowledges support from the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq.

REFERENCES

1. D. Hadjimichef, G. Krein, S. Szpigel and J. S. da Veiga, Ann. of Phys. **268** 105 (1998).
2. E. S. Ackleh, T. Barnes and E. S. Swanson, Phys. Rev. D **54** 6811 (1996).